

COGUMELOS NA BEIRA INTERIOR



Agaricus arvensis (Bola de neve)



Agaricus bisporus (Agárico dos jardins)




Agaricus campestris (Rosa dos prados)



Lactarius deliciosus (Sancha)



Agaricus sylvicola (Agárico da floresta)



Amanita caesarea (Ovo de rei)



Amanita citrina (Miscaro limão)



Amanita muscaria (Mata moscas)



Macrolepiota procera (Frade)



Amanita pantherina (Pantera)



Amanita ponderosa (Tortulho)



Amanita rubescens (Pé vermelho)



Armillaria mellea (Tarlhico)



Boletus aereus (Boleto escuro)



Boletus pinophilus (Boleto do pinheiro)



Cantharellus cibarius (Cantarelo)



Tricholoma equestre (Miscaro amarelo)



Coprinus comatus (Cabeludo)



Fistulina hepatica (Língua de vaca)



Hydnum repandum (Pata de borrego)



Lentinus edodes (Shitake)



Boletus edulis (Cépa)



Lepista nuda (Pé azul)



Pleurotus ostreatus (Repolga)



Russula sp. (Russula)



Suillus bellinii (Suito)



Terfezia arenaria (Criadilha)



Tricholoma portentosum (Miscaro branco)

Autor: José Luís Gravito Henriques, Eng. Agrónomo
Com base em Inquérito